

References

1. U.S. Department of Health and Human Services. Reducing Tobacco Use: A Report of the Surgeon General-Executive Summary. Atlanta, Georgia: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2000.
2. Cigarette Smoking Among Adults — United States, 2000, *MMWR* Highlights, July 26, 2002 / Vol. 51 / No. 29
9. Kottke, T.E., Battista, R.N., DeFries, G.H., et al. (1988). Attributes of successful smoking cessation intervention in medical practice: a meta-analysis of 39 controlled trials. *JAMA*; 259:2882-2889.
10. Cummings, S.R., Coates, T.J., Richard, R.J., et al. (1989). Training physicians in counseling about smoking cessation: a randomized trial of the 'Quit for Life' program. *Ann Intern Med*; 110:640-647.
11. Strecher, V.J., O'Malley, M.S., Villagra, V.G., et al. (1991). Can residents be trained to counsel patients about quitting smoking? Results from a randomized trial. *J Gen Intern Med*; 6: 9-17.
12. Schwartz, J.L. (1987). Review and evaluation of smoking cessation methods: the United States and Canada. Washington, DC: United States Department of Health and Human Services. National Institutes of Health Publication 87-2940
13. Cromwell, J., Bartosch, W.J., Fiore, M.C., et al. (1997). Cost-effectiveness of the clinical practice recommendations in the AHCPR guideline for smoking cessation. *JAMA*; 278:1759-1766.
14. Jamrozik, K., Vessey, M., Fowler, G., et al. (1984). Controlled trial of three different anti-smoking interventions in general practice. *BMJ*; 288:1499-1503.
15. Richmond, R.L., and Webster, I.W. (1985). A smoking cessation programme for general practice. *Med J Aust*; 142:190-194.
16. Risser, N.L., Belcher, D.W. (1990). Adding spirometry, carbon monoxide and pulmonary symptom results to smoking cessation counseling: a randomized trial. *J Gen Intern Med*; 5:16-22.
16. Lando, H.A., McGovern, P.G., Barrios, F.X., et al. (1990). Comparative evaluation of American Cancer Society and American Lung Association smoking cessation clinics. *Am J Public Health*; 80:554-559.
17. Thompson RS, Michnich ME, Friedlander L, et al. (1988). Effectiveness of smoking cessation interventions integrated into primary care practice. *Med Care*; 26:62-76.

18. Strecher, V.J., Krauter, M., Den Boer, D.J., et al. (1994). The effects of computer tailored smoking cessation messages in family practice settings. *J Fam Pract*; 39:262-270.
19. Prochaska, J.O., DiClemente, C.C., Velicer, W.F., Rossi, J.S. (1993). Standardized, individualized, interactive, and personalized self-help programs for smoking cessation. *Health Psychol*; 12:399-405.
20. Velicer, W.F., Prochaska, J.O. (1999). An expert system intervention for smoking cessation. *Pt Educ & Counseling*; 36:119-129.
21. Velicer, W.F., Prochaska, J.O., Bellis, J.M., DiClemente, C.C. et al. (1993). An expert system intervention for smoking cessation. *Addict Beh*; 18:269-290.
22. Velicer, W.F., Prochaska, J.O., Fava, J.L., LaForge, R.G., Rossi, J.S. (1999). *Interactive versus noninteractive interventions and dose-response relationships for stage-matched smoking cessation programs in a managed care setting.* *Health Psychol*; 18:21-28.
23. O'Neill, H.K., Gillispie, M.A., Slobin, K. (2000). Stages of change and smoking cessation: A computer-administered intervention program for young adults. *Am J Health Promotion*, 15:93-96.
24. Pallonen, U.E., Leskinen, L., Prochaska, J.O., Willey, C.J., et al. (1994). A 2-year self-help smoking cessation manual intervention among middle-aged Finnish men: An application of the transtheoretical model. *Prev Med*, 23:507-514.
25. Lennox A.S., Osman, L.M., Reiter, E., Robertson, R., et al. (2001). Cost effectiveness of computer tailored and non-tailored smoking cessation letters in general practice: Randomised controlled trial. *BMJ*; 322(7299):1396-1400.
26. Edwards, N.B., Simmons, R.C., Rosenthal, T.L., et al. (1988). Doxepin in the treatment of nicotine withdrawal. *Psychosomatics*; 29:203-206.
30. Fiscella, K., Franks, P. (1996). Cost-effectiveness of the transdermal nicotine patch as an adjunct to physicians' smoking cessation counseling. *JAMA*, 275:1247-1251.
31. Orleans, C.T., Resch, N., Noll, E., et al. (1994). Use of transdermal nicotine in a state-level prescription plan for the elderly: a first look at "real-world" patch users. *JAMA*; 271:601-607.
32. Fiore, M.C., Smith, S.S., Jorenby, D.E., et al. (1994). The effectiveness of nicotine patch for smoking cessation: a meta-analysis. *JAMA*; 271:1940-1947.
33. Hurd, R.D., Sachs, D.P.L., Glover, E.D., et al. (1997). A comparison of sustained release bupropion and placebo for smoking cessation. *N Engl J Med*; 337:1195-1202.

34. Russell, M.A.H., Stapelton, J.A., Hajek, P., et al. (1988). District programme to reduce smoking: can sustained intervention by general practitioners affect prevalence? *J Epidemiol Community Health*; 42:111-115.
35. Solberg, L.I., Maxwell, P.L., Kottke, T.E., et al. (1990). A systematic primary care office-based smoking cessation program. *J Fam Pract*; 30:647-654.
37. McKenna, K., Higgins, H. (1997). Factors influencing smoking cessation in patients with coronary artery disease. *Pt Educ & Counseling*; 32:197-205.
38. Murray, R.P., Gerald, L.B., Lindgren, P.G., Connett, J.E., et al. (2000). Characteristics of participants who stop smoking and sustain abstinence for 1 and 5 years in the Lung Health Study. *Prev Med*; 30:392-400.
39. Sanders, D., Peveler, R., Mant, D., Fowler, G. (1993). Predictors of successful smoking cessation following advice from nurses in general practice. *Addiction*; 88:1699-1705.
40. Wetter, D.W., Kenford, S.L., Smith, S.S., Fiore, M.C., et al. (1999). Gender differences in smoking cessation. *JCCP*; 67:555-562.
41. Matheny, K.B., Weatherman, K.E. (1988). Predictors of smoking cessation and maintenance. *J Clinical Psychol*; 54:223-235.
42. Klesges, R.C., Haddock, C.K., Lando, H., Talcott, G.W. (1999). Efficacy of forced smoking cessation and an adjunctive behavioral treatment on long-term smoking rates. *JCCP*; 67:952-958.
43. Glasgow, R.E., Klesges, R.C., O'Neill, H.K. (1986). Programming social support for smoking modification: An extension and replication. *Addict Beh*; 11:453-457.
44. Schoenbach, V.J., Orleans, C.T., Wagner, E.H., Quade, D., et al. (1992). Characteristics of smokers who enroll and quit in self-help programs. *Health Educ Res*; 7:369-380.
56. Fiore, M.C. (2000). A clinical practice guideline for treating tobacco use and dependence: A US Public Health Service report. *JAMA*; 283:3244-3254.